

# DENTISTRY TODAY

Circulation 150,000

VOLUME 28 NO. 8

THE NATION'S LEADING CLINICAL NEWS MAGAZINE FOR DENTISTS

AUGUST 2009

## IMPLANTS



# Optimizing Implant Placement and Aesthetics: Technology to the Rescue!



Timothy  
Kosinski, DDS

The continually evolving science of implant dentistry has led to a growing recognition that many treatment options offer predictable long-term results. Dental implants have certainly developed into a viable alternative to conventional prosthodontic reconstruction of edentulous spaces, and have afforded us a restorative tool for use in difficult aesthetic cases. Optimizing aesthetics in the anterior maxillary incisor area is critical to a successful outcome.

The therapeutic goal of implant dentistry is not merely tooth replacement, but total oral rehabilitation. Implants provide excellent support for fixed appliances, increasing function as compared to conventional dental therapies. Implant dentistry has gone through many phases over the years. Modern design allows us to predictably place our dental implants in immediate extraction sites.<sup>1-3</sup> Single tooth-by-tooth reconstruction provides easy access for the patient to floss and clean the areas, compared with



Figures 1 and 2. NobelGuide (Nobel Biocare) software used to idealize implant size and position.

the relative difficulty in maintenance when crowns are splinted. There is now little risk of abutments loosening under function, as was the case in the past.

This case study will serve to demonstrate a maxillary anterior reconstruction using immediate placement of dental implants, following diagnosis using computed tomography (CT) scanning software, immediate loading using

transitional abutments, splinted composite crowns, and dynamic and aesthetic smile design using zirconia abutments and free standing all-ceramic crowns.<sup>4-6</sup>

### ADVANTAGES OF COMPUTED TOMOGRAPHY SCANNING TECHNOLOGY

CT scanning is becoming more and more prevalent for use in ensuring proper dental implant

placement. Bitewings, periapicals, panoramic radiographs, and even medical CT scans only give a 2-dimensional (2-D) image for implant planning. One of the latest CT planning software (NobelGuide CT [Nobel Biocare]) provides as much information as possible to assist during surgical planning. We can now visualize vital anatomy in 2-D and 3-dimensional (3-D) prior to surgery, and can assess the location of implants virtually prior to any surgical intervention.<sup>7</sup> Diagnosing any anatomic issues specific to the patient, planning the implant type, determining position and orientation in the bone, are all easily accomplished with the help of this technology. The process is both user-friendly and intuitive.

The NobelGuide CT scanning software as demonstrated below allows us to simulate the placement of implants accurately before ever touching the patient. A surgical guide, created from the 3-D images, helps place the implants in the proper positions, without the need for a flap incision.<sup>8,9</sup> This technique proves to be a cost effective solution to assist the implant dentist in planning an aesthetic final result and minimizing any surgical challenges that are faced. The CAD/CAM planning and placement system provided by NobelGuide provides a high level of comfort and safety for the patient by reducing surgical and restorative time. This is done by utilizing an accurate 3-D plan prior to implant placement.

There are obvious advantages including; easy visual understanding for clear case presentations, reduced surgical chair time, reduced restorative chair time in certain cases, reduced stress for the clinician and the patient, the avoidance of surprises during surgery, implants that are placed optimally for long-term implant and prosthetic success and, most importantly, an improved aesthetic result.<sup>10-12</sup>

### COMPUTED TOMOGRAPHY SCANNING TECHNIQUE

Prior to the CT scan, a radiographic guide is fabricated. This will serve to aid in the visualization of the optimal prosthetic outcome. The teeth are positioned properly in wax, and then a hard model



Figure 3. Preoperative dentition, retracted view.

is made to illustrate what the case will look like finished before ever starting. All appropriate dental anatomy is included. The radiographic guide is placed in the mouth during the CT scan. This allows us to see the ideal position of the teeth on a 3-D model. The entire 3-D image is analyzed and the implant planning and simulation of implant placement completed using the computer (Figures 1 and 2). The surgical placement of dental implants can be done in a conventional manner using the newly created surgical guide to help direct the implant in the ideal position. However, optimally the surgery can be completed without making any incisional flap. The implants are placed to the desired depth using the computer software and surgical guide.

### MATERIALS AND METHODS: ADVANTAGES OF TAPERED DESIGN

The Replace Implant System (Nobel Biocare) has evolved considerably since its introduction in 1997. The implants and all related surgical and prosthetic components incorporate a color coded system that allow users to identify at a glance which restorative parts go with which size implant placed. In 1999, the Replace Tapered Implants became available with an internal connection that simplifies the impression technique, seating of abutments and crown and bridge placement. Three locking channels guide the positioning of the abutment. Once torqued into place, the zirconia abutment does not loosen making single tooth restoration reliable. The system can be used in one or 2 stage surgical procedures. Primary stability is the key factor for successful early and immediate loading.

The tapered design of the Replace



Figure 4. Radiograph illustrating severe root resorption.


Select gives placement alternatives in sites with anatomical limitations, such as labial concavities in the pre-maxilla and converging adjacent tooth roots. Since they approximate the shape of a natural tooth root, they provided better stability in extraction sites. There are several collar heights and designs available in these titanium surface-treated (TiUnite [Nobel Biocare]) implants; including no collar, 1.5 mm and 2.0 mm. The shorter collar designs are indicated for aesthetic areas. Nobel Biocare's Procera process produces custom-shaped zirconium oxide abutments to create the most natural form and emergence profile. The ultimate aesthetic solution, especially for patients with a high smile line and thin tissue, is achieved using the Procera abutments. Every aspect of the implant system makes the restoration of teeth on implants as easy as crown and bridge.<sup>13-15</sup>

The Replace Select implant has a tapered body and an internal prosthetic connection. The internal connection is trichanneled. The internal lengths of the channels are approximately 1.5 mm. The implant has a wide crestal interface with a tapered thread design. The diameters, at the crest of the implant, are 3.5 mm, 4.3 mm, 5.0 mm, and 6.0 mm. This allows for a better emergence profile from the top of the implant. Anatomical considerations for the use of the implant





**Figures 5 and 6.** Laboratory casts and components used for immediate transitional splinted crowns for teeth Nos. 7 to 10.



**Figure 7.** Flapless extraction site.



**Figures 8 and 9.** NobelGuide surgical stent with precision openings for drills.



**Figure 10.** Surgical stent in place with narrow platform guide, used for pilot drill angulation and depth.



**Figure 11.** A 3.5-mm surgical drill.



**Figure 12.** A 3.5-mm Replace Select (Nobel Biocare) dental implant treated to depth and angulation using the surgical guide.



**Figures 13.** A 4.3-mm Replace Select dental implants positioned.

include converging roots and concavities with the bone, which makes placing a parallel-walled implant more difficult. Interface of the implant increases dramatically as implant diameter increases. This may be more relevant than implant length.

The tapered design of the implant promotes elevated levels of fatigue endurance since the coronal portion is wider in diameter than the apical portion. This taper also ensures a tight fit and promotes function coronally to help offset stress shielding along the narrow, smooth crestal band on the implant. This reduces bone resorption that may result from hypo function. This is especially important during placement of

the implant in a fresh extraction site.

The taper also reduces the incidence of cortical plate bone perforation during osteotomy preparation near anatomic undercut areas and protects adjacent natural tooth roots. The tapered design of the implant also often allows for better angulation of the implant. It is important to have an implant design that allows for placement of the fixture in a way that places the forces down the long axis. The implant has versatility and allows for aesthetic and variable bone morphology.


#### CASE REPORT

A 38-year-old female presented with mobile maxillary anterior lateral and


central incisors. These teeth had been orthodontically treated in the years before resulting in resorption, mobility, and aesthetic problems (Figure 3). Her main concerns were that she would have to go without teeth for any length of time, or the possibility of having to wear a removable appliance. The patient requested something she referred to as, "teeth in a day." Oral and radiographic evaluation indicated severe root resorption around teeth Nos. 7 to 10 (Figure 4). Her teeth were deemed to be untreatable using traditional dental techniques.

#### Diagnosis and Treatment Planning

The decision was made to have a cone beam CT scan done and evaluated using



**Figures 14.** Implants placed ideally with flapless procedure.



**Figure 15.** Immediate conical abutments positioned.




**Figures 16.** Retracted view of transitional splinted crowns, cemented temporarily during healing.



**Figure 17.** Procera abutments torque into position.



**Figure 18.** Individual Procera crowns in place.



**Figure 19.** Panoramic view of implant, zirconia abutments, and crowns in place.

the NobelGuide CT scanning software. Virtually placing the dental implants using the computer, prior to ever touching the patient, would prove to be a tremendous advantage in diagnosing and preparing for this case (Figures 5 and 6).

There were no medical conditions, allergies, or sensitivities that would preclude the use of necessary implant procedures or medications. Since there was no need to work around any habits involving the use of nicotine or alcohol, healing was expected to be good and uncomplicated.

Proper diagnosis for dental reconstruction is the most critical aspect of any surgical intervention. Determining the design of the final prosthetic reconstruction may be the most difficult procedure. It is this author's belief that *the restoring dentist should be dictating dental implant position and angulation prior to surgical placement.* To help achieve this goal, diagnostic wax-ups, modern computer generated scanning, or simple common-sense design should be considered. The diagnostic wax-up is an invaluable aid in determining the proper placement of implants in the center, between the proposed abutment teeth. The wax-

up is particularly valuable when cement-retained or screw-retained crown and bridge is planned. NobelGuide CT scanning software is used to create a surgical guide that is used pre-surgically in determining the ideal location and angulation of the implant site, and to optimize placement of dental implants for maximum aesthetic and functional results.

#### **Surgical Stent, Extractions, and Implant Placement**

Prior to extraction of the mobile teeth (Figure 7) a surgical stent was fabricated using a master cast of her existing bridgework (Figures 8 and 9). Tapered drills of increasing widths were used to prepare the bone to accept the proper size implant. 3.5 x 13 mm Replace Select dental implants were guided into place using the NobelGuide surgical stent in the Nos. 7 and 10 area and 4.3 x 13 mm Replace Select implants surgically guided into the Nos. 8 and 9 areas (Figures 10 and 11).

A color-coded threadformer, corresponding to the implant diameter selected, was then used. The maximum recommended setting is 30 rpm. Firm pressure was applied to the threadformer

and it was rotated slowly. Once the threads were engaged, the threadformer was allowed to feed without pressure. The osteotomy was threaded to the single depth reference line on each drill. The dental implants were then removed from their sterile packaging and threaded into the prepared site. (When more torque is needed to complete the placement, a ratchet and insertion assembly is used to place the implant to its final depth. A marking indicates that one of the 3 trichanneled internal connections is placed to the facial.) (Figures 12 to 14.)

Immediate loading of dental implants have proven to be a predictable method of improving emergence profile and smile design. The Nobel Biocare Immediate temporary abutments were used to secure a transitional splinted crowns and allow the gingiva to respond positively (Figure 15). Immediate aesthetics and function were achieved (Figure 16). Simple color-coded transfer assembly can duplicate the position of the dental implant on a working cast. Procera abutments were reshaped in the dental laboratory with slightly subgingival margins. These abutments were tightened to 35 Ncm to ensure that they would not loosen (Figure 17).





**Figure 20.** Final smile line, with individual all-ceramic crowns (Procera Zirconia [Nobel Biocare]) in place.

### SUMMARY

The goal for this patient was to create an aesthetic smile design using individual dental implants to reconstruct the edentulous spaces. Procera crowns aesthetically restored the separate and distinct teeth (Figures 18 and 19). The patient was thrilled with the final aesthetic result, and she was able to smile confidently again (Figure 20).◆

### References

1. Locante WM. Single-tooth replacements in the esthetic zone with an immediate function implant: a preliminary report. *J Oral Implantol.* 2004;30:369-375.
2. Degidi M, Piattelli A. Comparative analysis study of 702 dental implants subjected to immediate functional loading and immediate nonfunctional loading to traditional healing periods with a follow-up of up to 24 months. *Int J Oral Maxillofac Implants.* 2005;20:99-107.
3. Villa R, Rangert B. Early loading of interforaminal implants immediately installed after extraction of teeth presenting endodontic and periodontal lesions. *Clin Implant Dent Relat Res.* 2005;7(suppl 1):28-35.
4. Glauser R, Rée A, Lundgren A, et al. Immediate occlusal loading of Brånemark implants applied in various jawbone regions: a prospective, 1-year clinical study. *Clin Implant Dent Relat Res.* 2001;3:204-213.
5. Lozada JL, Ardah AJ, Rungcharassaeng K, et al. Immediate functional load of mandibular implant overdentures: a surgical and prosthodontic rationale of 2 implant modalities. *J Oral Implantol.* 2004;30:297-306.
6. Ganeles J, Wismeijer D. Early and immediately restored and loaded dental implants for single-tooth and partial-arch applications. *Int J Oral Maxillofac Implants.* 2004;19(suppl):92-102.
7. van Steenberghe D, Glauser R, Blombäck U, et al. A computed tomographic scan-derived customized surgical template and fixed prosthesis for flapless surgery and immediate loading of implants in fully edentulous maxillae: a prospective multicenter study. *Clin Implant Dent Relat Res.* 2005;7(suppl 1):111-120.
8. Balshi SF, Wolfinger GJ, Balshi TJ. A prospective study of immediate functional loading, following the Teeth in a Day protocol: a case series of 55 consecutive edentulous maxillas. *Clin Implant Dent Relat Res.* 2005;7:24-31.
9. Maló P, Rangert B, Nobre M. All-on-4 immediate-function concept with Brånemark System implants for completely edentulous maxillae: a 1-year retrospective clinical study. *Clin Implant Dent Relat Res.* 2005;7(suppl 1):88-94.
10. Lee CY. Immediate load protocol for anterior maxilla with cortical bone from mandibular ramus. *Implant Dent.* 2006;15:153-159.
11. Olsson M, Urde G, Andersen JB, et al. Early loading of maxillary fixed cross-arch dental prostheses supported by six or eight oxidized titanium implants: results after 1 year of loading, case series. *Clin Implant Dent Relat Res.* 2003;5(suppl 1):81-87.
12. Ostman PO, Hellman M, Sennerby L. Direct implant loading in the edentulous maxilla using a bone density-adapted surgical protocol and primary implant stability criteria for inclusion. *Clin Implant Dent Relat Res.* 2005;7(suppl 1):60-69.
13. Scortecchi G. Immediate function of cortically anchored disk-design implants without bone augmentation in moderately to severely resorbed completely edentulous maxillae. *J Oral Implantol.* 1999;25:70-79.
14. Babbush CA. Provisional implants: surgical and prosthetic aspects. *Implant Dent.* 2001;10:113-120.
15. Becker W, Wong J. Early functional loading in the fully edentulous mandible after mandibular resection and reconstruction due to an ameloblastoma: case report. *Clin Implant Dent Relat Res.* 2003;5:47-51.

**Dr. Kosinski** is an adjunct assistant professor at the University of Detroit Mercy School of Dentistry. He received his DDS from the University of Detroit Mercy Dental School and his Mastership in Biochemistry from Wayne State University School of Medicine. He is a Diplomate of the American Board of Oral Implantology/Implant Dentistry, the International Congress of Oral Implantologists and the American Society of Osseointegration. He is a Fellow of the American Academy of Implant Dentistry and received his Mastership in the Academy of General Dentistry. Dr. Kosinski serves on the editorial review board of *REALITY*, the information source for aesthetic dentistry, *Contemporary Esthetics and Clinical Advisors*, and became the editor of the Michigan Academy of General Dentistry. Dr. Kosinski has received many honors including Fellowship in the American and International Colleges of Dentists and the Academy of Dentistry International. He is a member of OKU and the Pierre Fauchard Academy. Dr. Kosinski has published over 50 articles on the surgical and prosthetic phases of implant dentistry and he was a contributor to the textbook *Principles and Practices of Implant Dentistry*. He can be reached via e-mail at allquestions@smilecreator.net.

*Disclosure:* Dr. Kosinski reports no conflict of interest.

**SMILECREATOR of  
Bingham Farms<sup>SM</sup>**

Timothy Kosinski, DDS  
31000 Telegraph Rd., Suite 170  
Bingham Farms, MI 48025  
248-646-8651 • fax 248-646-8659  
allquestions@smilecreator.net  
www.smilecreator.net

PROOF COPY ONLY

WWW.SMILECREATOR.NET

Dental Implants \* Comprehensive Dentistry \* Smile Design \* Care for the Apprehensive Patient